

PPA and the Risk of Hemorrhagic Stroke

Lewis H. Kuller, M.D., Dr.P.H., M.P.H.
Chairman of Epidemiology
University of Pittsburgh School of Health

9

PPA and the Risk of Hemorrhagic Stroke Selection Bias

- Basic problems with case and control selection in the conduct of the study render this a failed study.
- Cases
 - 702 cases out of 1717 potential cases (41%)
- Controls
 - 151 telephone numbers called to identify a control (p. 17, final report)
 - Approximately 3 eligibles for each control

10

**Cases and controls differ greatly on
demographic characteristics,
suggesting very substantial selection bias**

<u>Education</u>	<u>Cases (%)</u>	<u>Controls (%)</u>
College	40	62
< High School	20	9

11

Use History

Any PPA Exposure

Cases -- 3.8 %

Controls -- 2.4%

- Assume that prevalence was 4% in the 4200 controls not included. True prevalence in controls would be 3.6%

$$1400 \times 2.4 = 33$$

$$4200 \times .04 = 168$$

- Same as cases: Due to select bias of controls -- unknown true prevalence of PPA in controls

12

Time Period of Use of Drug (Men and Women)

	Cases		Controls	
	N	%	N	%
Within 2 wks of current use				
First use	8	1.1	5	0.4
Not first use	13	1.9	16	1.2
Prior use	06	0.9	12	0.9
<i>Use 2 wks to 3 days of event</i>	11	1.7	33	2.5
Total use	38	5.4	66	4.8

Why greater use in controls -2 weeks-3days prior to event?

13

Sex Differences in Outcome

Any PPA <3 days	<u>Cases (n=383)</u>		<u>Controls (n=750)</u>		<u>O.R.</u>
	N	%	N	%	
Women	21	5.5	20	2.7	2.0
	<u>Cases (n=319)</u>		<u>Controls (n=626)</u>		
	N	%	N	%	
Men	6	1.9	13	2.1	-

Difference in women cases only

14

Cough/Cold Only

Any PPA <3 days	<u>Cases</u>		<u>Controls</u>		<u>O.R.</u>
	N	%	N	%	
Women	16	4.2	19	2.5	1.54
Men	6	1.9	13	2.1	0.62

Risk for women, protective for men?

15

Small Numbers

- Appetite Suppressant for Women

6 Cases
1 Control
Odds Ratio = 12

- Prior Use in Men

1 Case
8 Control
Odds Ratio = 0.25

16

Characteristics of Cases

- Consumption of Caffeine:
Current Users (first use in 3 days)
 - Cases (9)
 - 10 cups coffee / day
 - 8.5 cups coffee / day
 - 10 glasses soda / day
 - 8 glasses soda / day
 - 6 glasses soda / wk (History of stroke)
 - 1 glass soda / wk (History of stroke)
 - 2 - (0) (Both prior headaches)
 - Controls (5)
 - 6 cups / day coffee
 - 6 glasses / day soda
 - 2 cups coffee / day (2)
 - 1 negative / day (cup of tea)

17

18